

Week 6 Worksheet

Electrostatic Energy

Jacob Erlichman

Exercise 1. In this exercise, you will find the energy contained in the electric field \mathbf{E} .

- How much work does it take to assemble a configuration of two point charges?
- What about n point charges?
- Show that your result from (b) can be written

$$W = \frac{1}{2} \sum_{i=1}^n q_i V(\mathbf{r}_i),$$

where q_i is the charge of the i^{th} charge and \mathbf{r}_i is its position vector.

- Generalize your result from (c) to a *continuous* charge distribution with (not necessarily uniform) charge density ρ .
- Using the differential form of Gauss' law, $\rho = \epsilon_0 \nabla \cdot \mathbf{E}$, rewrite this in terms of \mathbf{E} and V . Integrate by parts to transfer the derivative ∇ to V instead of \mathbf{E} , and argue that your answer is exactly

$$W = \frac{\epsilon_0}{2} \int E^2 dV,$$

where the integral is taken over *all space*.

Hint: Integration by parts in this context can be achieved by using the product rule on $\nabla \cdot (\mathbf{v} f)$ (also the divergence theorem will help). Notice the similarity to 1-dimensional integration by parts, which can be derived from the identity $\frac{d}{dx}(fg) = f'g + fg'$.

Exercise 2. Griffiths 2.39. Two spherical cavities of radii a and b , respectively, are hollowed out from the interior of a neutral conducting sphere of radius R . At the center of each cavity is a point charge, q_a and q_b , respectively.

- Find the surface charge densities σ_a , σ_b , and σ_R .
- What is the field outside the conductor?
- What is the field within each cavity?
- What is the force on q_a and q_b ?
- Use the previous exercise to find the energy of this configuration.