

Week 5 Worksheet Solutions

Symmetries!

Jacob Erlikhman

September 30, 2024

Exercise 1. In this problem, you will construct the 2×2 matrix corresponding to a finite rotation which places the \hat{z} axis along an arbitrary direction \hat{r} .

- a) A rotation can be specified by the Euler angles (α, β, γ) , or by (θ, φ) . The Euler angles represent first a rotation about \hat{z} by an angle α , then a rotation *about the new* y -axis by an angle β , and then a rotation about the *new* z -axis again. Convince yourself that this works.
- b) Now, suppose given a rotation specified by the Euler angles (α, β, γ) . This is given in quantum mechanics by the matrix

$$e^{-i\gamma S_{z'}/\hbar} e^{-i\beta S_u/\hbar} e^{-i\alpha S_z/\hbar},$$

where the u -axis is the new y -axis after rotating about z , and the z' -axis is the new z -axis after rotating about \hat{z} and \hat{u} . Show that this is the same matrix as

$$e^{-i\alpha S_z/\hbar} e^{-i\beta S_y/\hbar} e^{-i\gamma S_z/\hbar}.$$

Hint: Denoting a rotation about the axis r by an angle ζ as $R_r(\zeta)$, we have that $S_u = R_z(\alpha)S_yR_z(-\alpha) = e^{-i\alpha S_z/\hbar}S_ye^{i\alpha S_z/\hbar}$. Now, try to write a similar expression for $R_{z'}(\gamma) = e^{-i\gamma S_{z'}/\hbar}$.

- c) Use part (b) with $S_i = \frac{\hbar}{2}\sigma_i$ to calculate the rotation matrix corresponding to placing the \hat{z} axis along \hat{r} , where \hat{r} is specified by the two angles (θ, φ) .
Hints: The idea is to Taylor expand each exponential. Think about a simple expression for σ_i^n , where σ_i is the Pauli matrix you need. Finally, one of the results you should get along the way is

$$e^{-i\beta\sigma_y/2} = \cos(\beta/2)\mathbb{1} - i\sigma_y \sin(\beta/2).$$

- d) **Griffiths 6.32(f).** Calculate the matrix corresponding to a rotation by π about \hat{x} .
- e) **Griffiths 6.32(g).** Calculate the matrix corresponding to a 2π rotation about \hat{z} . Comment on the answer.
- a) Convinced!

b) We have $S_u = R_z(\alpha)S_yR_z(-\alpha)$ and $S_{z'} = R_u(\beta)S_zR_u(-\beta)$. First of all, note that

$$\begin{aligned} R_u(\beta) &= \exp(-i\beta R_z(\alpha)S_yR_z(-\alpha)) \\ &= \sum_{n=0}^{\infty} \frac{(-i\beta)^n}{n!} (R_z(\alpha)S_yR_z(-\alpha))^n. \end{aligned}$$

Now, observe that $(R_z(\alpha)S_yR_z(-\alpha))^n = R_z(\alpha)S_y^nR_z(-\alpha)$. Thus,

$$\begin{aligned} R_u(\beta) &= R_z(\alpha) \sum_{n=0}^{\infty} \frac{(-i\beta S_y)^n}{n!} R_z(-\alpha) \\ &= R_z(\alpha) e^{-i\beta S_y} R_z(-\alpha) \\ &= R_z(\alpha) R_y(\beta) R_z(-\alpha). \end{aligned}$$

Likewise,

$$R_{z'}(\gamma) = R_u(\beta)R_z(\gamma)R_u(-\beta).$$

Putting everything together, we find

$$R_{z'}(\gamma)R_u(\beta)R_z(\alpha) = R_z(\alpha)R_y(\beta)R_z(\gamma).$$

c) We actually only need two Euler angles to achieve this, α and β , with $\theta = \beta$ and $\varphi = \alpha$. So we get

$$e^{-i\varphi\sigma_z/2}e^{-i\theta\sigma_y/2}.$$

We work one term at a time. The first term is easy since σ_z is diagonal (recall [or immediately prove!] that for a diagonal matrix $D = (d_1, \dots, d_n)$, $e^D = (e^{d_1}, \dots, e^{d_n})$), so

$$e^{-i\varphi\sigma_z/2} = \begin{bmatrix} e^{-i\varphi/2} & 0 \\ 0 & e^{i\varphi/2} \end{bmatrix}.$$

For the second term, we get

$$e^{-i\theta\sigma_y/2} = \mathbb{1} \cos(\theta/2) - i\sigma_y \sin(\theta/2).$$

If you don't see this right away, try to write out the power series expansion, remembering that $\sigma_y^2 = \mathbb{1}$. Explicitly, we have

$$e^{-i\theta\sigma_y/2} = \sum_{j=0}^{\infty} \left(\frac{-i\theta}{2}\right)^{2j} \frac{1}{(2j)!} \mathbb{1} + \sum_{k=0}^{\infty} \left(\frac{-i\theta}{2}\right)^{2k+1} \frac{1}{(2k+1)!} \sigma_y.$$

Now, $(-i)^{2j} = (-1)^j$, while $(-i)^{2k+1} = -i(-1)^k$. Hence, the first term can be recognized as the power series expansion for $\cos(\theta/2)$, and the second as the power series expansion for $-i \sin(\theta/2)$. Putting it all together, we find

$$e^{-i\varphi\sigma_z/2}e^{-i\theta\sigma_y/2} = \begin{bmatrix} e^{-i\varphi/2} \cos(\theta/2) & -e^{-i\varphi/2} \sin(\theta/2) \\ e^{i\varphi/2} \sin(\theta/2) & e^{i\varphi/2} \cos(\theta/2) \end{bmatrix}.$$

- d) This is just $e^{-i\pi\sigma_x/2}$. Notice that $\sigma_x^2 = \mathbb{1}$, so that $e^{-i\pi\sigma_x/2} = \cos(\pi/2)\mathbb{1} - i \sin(\pi/2)\sigma_x = -i\sigma_x$.
- e) This is $e^{-i\pi\sigma_z} = -\mathbb{1}$. Thus, a 2π rotation about \hat{z} of a spin 1/2 particle returns *negative* the particle state! This is a purely quantum mechanical phenomenon (and can be measured in practice).

Exercise 2. Another symmetry is called **dilation** symmetry. Dilations are given by the transformation $\mathbf{x} \rightarrow \mathbf{x}' = e^c \mathbf{x}$, where $c \in \mathbb{R}$. Call its generator D , so that e^{-icD} is the corresponding unitary operator.

- a) Show that the *infinitesimal* transformation

$$e^{i\mathbf{a}\cdot\mathbf{p}} e^{icD} e^{-i\mathbf{a}\cdot\mathbf{p}} e^{-icD}$$

is given by $\mathbb{1} + c\mathbf{a} \cdot [D, \mathbf{p}]$.

- b) Calculate $[D, \mathbf{p}]$.

This problem will be on the midterm review, so solutions will be posted along with solutions to the midterm review problems.