## Week 5 Worksheet Solutions More Perturbation Theory

Jacob Erlikhman

October 1, 2025

Exercise 1. Griffiths 7.54. Last week, you derived the first order correction to the expectation value of an observable A in the n<sup>th</sup> energy eigenstate of a system perturbed by H'. You found

$$\langle A \rangle^1 = 2 \sum_{m \neq n} \frac{\left\langle \psi_n^0 | A | \psi_m^0 \right\rangle \left\langle \psi_m^0 | H' | \psi_n^0 \right\rangle}{E_n^0 - E_m^0}.$$

Suppose we have a particle of charge q in a weak electric field  $\mathbf{E} = E_{\rm ext}\hat{x}$ , so that  $H' = -qE_{\rm ext}x$ . This induces a dipole moment  $p_e = qx$  in the "atom." The expectation value of  $p_e$  is proportional to the applied field, and the proportionality factor is called the **polarizability**,  $\alpha$ . Show that

$$\alpha = -2q^2 \sum_{m \neq n} \frac{\left| \left\langle \psi_n^0 | x | \psi_m^0 \right\rangle \right|^2}{E_n^0 - E_m^0}.$$

Find  $\alpha$  for the ground state of a 1-D harmonic oscillator, and compare the classical answer.

*Hint:* Recall that x can be written in terms of creation and annihilation operators. Given

$$H^0 = \frac{1}{2m} \left[ p^2 + (m\omega x)^2 \right],$$

you can derive what a and  $a^{\dagger}$  should be in terms of x and p by using the sum of squares formula. To get the "usual" form, rescale each of them by  $a \to \frac{1}{\sqrt{\hbar\omega}}a$  (so that the hamiltonian can be written  $H^0 = a^{\dagger}a + 1/2$ ).

Just plug in A = qx into the formula for  $\langle A \rangle^1$ . Since  $H' = -qE_{\rm ext}x$ , we get exactly the formula for  $\alpha$  shown. Now, we want to find  $\alpha$  for the ground state of the harmonic oscillator. In fact, we can find it for any energy eigenstate. Denote the  $n^{\rm th}$  energy eigenstate by  $|n\rangle$ , so that

$$\alpha = -2q^2 \sum_{m \neq n} \frac{|\langle n | x | m \rangle|^2}{(n-m)\hbar \omega}.$$

Worksheet 5 2

Following the derivation suggested in the hint, we get

$$a^{\dagger} = \frac{1}{\sqrt{2m}} (m\omega x - ip)$$
$$a = \frac{1}{\sqrt{2m}} (m\omega x + ip).$$

Now, rescale a and  $a^{\dagger}$  as indicated, so that

$$a^{\dagger} = \frac{1}{\sqrt{2m\hbar\omega}}(m\omega x + ip)$$
$$a = \frac{1}{\sqrt{2m\hbar\omega}}(m\omega x - ip).$$

Thus,

$$x = \sqrt{\frac{\hbar}{2m\omega}}(a + a^{\dagger}).$$

Plugging this in to the form for  $\alpha$  above, we get

$$\alpha = -\frac{\hbar q^2}{m\omega} \sum_{m \neq n} \frac{|\langle n|(a+a^{\dagger})|m\rangle|^2}{(n-m)\hbar\omega}$$
$$= -\frac{q^2}{m\omega^2} (n - (n+1))$$
$$= \frac{q^2}{m\omega^2}.$$

Classically, a charged particle in a harmonic oscillator potential with an electric field has equation of motion given by Newton's second law:

$$m\ddot{x} = -kx + qE_{\rm ext}.$$

The solution of this equation is just

$$x(t) = A\sin(\omega t) + B\cos(\omega t) + \frac{1}{k}qE_{\text{ext}},$$

where  $\omega = \sqrt{\frac{k}{m}}$ . Since the average value of a sinusoid is 0, we get  $\alpha = \frac{q^2}{m\omega^2}$ , which is exactly the same as the perturbation theory calculation! This is something that commonly occurs in perturbation theory calculations (especially in quantum field theory), where the first order correction just gives back the classical result. This necessitates going to second or higher order to get any truly quantum effects due to the perturbation!

Exercise 2. Griffiths 7.45. Stark Effect in Hydrogen. When an atom is placed in a uniform electric field  $\mathbf{E}_{\text{ext}}$ , the energy levels are shifted. This is known as the **Stark effect**. You'll analyze the Stark effect for the n=1 and n=2 states of hydrogen. Suppose  $\mathbf{E}_{\text{ext}} = E_{\text{ext}} \hat{z}$ , so that

$$H' = eE_{\rm ext}r\cos\theta$$

is the perturbation of the hamiltonian for the electron, where  $H^0 = \frac{p^2}{2m} - \frac{e^2}{4\pi\epsilon_0} \frac{1}{r}$ .

Worksheet 5 3

- a) Show that the ground state energy is unchanged at first order.
- b) How much degeneracy does the first excited state have? List the degenerate states.
- c) Determine the first-order corrections to the energy. Into how many levels does  $E_2$  split? Hint: All  $W_{ij}$  are 0 except for two, and you can avoid doing all of the zero integrals in this problem by using symmetry and selection rules. You'll need the following

$$\psi_{210} = \frac{1}{2\sqrt{6}} a^{-3/2} \frac{r}{a} e^{-r/2a} \sqrt{\frac{3}{4\pi}} \cos \theta$$

$$\psi_{200} = \frac{1}{\sqrt{2}} a^{-3/2} \left( 1 - \frac{r}{2a} \right) e^{-r/2a} \frac{1}{2\sqrt{\pi}}.$$

d) What are the "good" wavefunctions for (b)? Find the expectation value of the electric dipole moment in each of these states.

I think we'll have this problem on the midterm review, so solutions will be posted along with solutions to the review problems.